arXiv:gr-qc/9905034v2 24 Aug 1999

Black hole radiation with high frequency dispersion

Hiromi SAIDA[a‡] and Masa-aki SAKAGAMI[b§]
[a] *Graduate School of Human and Environmental Studies, Kyoto University, Kyoto 606-8501, Japan,*
[b] *Department of Fundamental Sciences, FIHS, Kyoto University, Kyoto 606-8501, Japan*

We consider one model of a black hole radiation, in which the equation of motion of a matter field is modified to cut off high frequency modes. The spectrum in the model has already been analytically derived in low frequency range, which has resulted in the Planckian distributin of the Hawking temperature. On the other hand, it has been numerically shown that its spectrum deviates from the thermal one in high frequency range. In this paper, we analytically derive the form of the deviation in the high frequency range. Our result can qualitatively explain the nature of the numerically calculated spectrum. The origin of the deviation is clarified by a simple discussion.

PACS number(s): 04.70.Dy, 04.62.+v, 04.70.-s

I. INTRODUCTION

There are two important theories in modern physics: quantum mechanics and the theory of general relativity. The former has been applied to field theory in flat spacetime and marked triumphs in understanding the wide range of microscopic phenomena. On the other hand, the latter has given us insights into phenomena having close relation to gravity. The theory of general relativity, which is a classical theory, has drastically changed our understandings of space and time. Yet, we have no idea to treat the gravity in Planck scale. It is unclear whether the unification of quantum mechanics and the theory of general relativity will be accomplished in the context of quantum physics. Today, however, we do not have any idea superior to the quantum mechanics in treating Planck scale physics. It is, therefore, meaningful to study how quantum and/or semi-classical effects will be expected in the presence of the strong gravity.

The most remarkable discovery including a semi-classical gravitational effect is the Hawking radiation [1], which is concluded by treating matter fields on spacetime as quantum while a black hole metric as classical. According to this theory, a black hole radiates particle flux of a thermal spectrum, whose temperature is $\kappa/2\pi$ where κ is the surface gravity.

The derivetion of the original Hawking radiation relies on the extremely high, over Planck scale, frequency modes. These modes arise from the extremely high gravitational red shift which the radiation undergoes during propagating from the event horizon to the asymptotically flat region. It is the investigation of the Hawking radiation that sheds light on Planck scale physics.

In the study of this physics, one model of a black hole radiation [3] [4] [5] has been proposed, which makes an assumption on the effects of unknown Planck scale physics as follows: the equation of motion (EOM) of matter field is modified to have a cut off of the high frequency modes with respect to free fall observer. [*] There are two important properties of this model. First one is that the modified EOM of matter field violates Lorentz invariance. Secondly, the origin of the black hole radiation we consider in this paper, which is called the mode conversion, [†] is different from that of the original Hawking radiation. We call the treatment of the black hole radiation in this paper as *the mode conversion* (*MC*) *model* hereafter.

The spectrum of the flux in two dimensional MC model has already been numerically calculated [3]. It has been shown that, although the spectrum almost agrees with the Planckian distributionits, it deviates from the thermal one in the high frequency region $\kappa \ll \omega$, where ω is the energy (frequency) of a massless scalar field. On the other hand, an analytical calculation [4] has resulted in the thermal spectrum, where the analysis has been carried out perturbatively up to lowest orders of κ and ω in the range $\omega < \kappa$. This is appropriate because the derivation of the spectrum in [4] does not take the effects of the high frequency range into account.

The purpose of this paper is to extend the analytical calculation in [4] to the high frequency range, $\kappa < \omega$. We obtain the form of the distribution function, $N(\omega)$, as

$$N(\omega) \simeq \frac{1 - \alpha\omega^2/2}{\exp[(2\pi\omega/\kappa)(1 - \alpha\omega^2/2)] - 1} ,$$

which shows the deviation from the thermal spectrum. Here, α is the square of the cut off scale characterizing the new physics in Planck scale. This result denotes the same tendency as what numerical calculations show. Furthermore, we will give a simple explanation of the occurrence of the deviation.

---

[*] This model was motivated by the dumb hole [2], which is a hydrodynamical analogue of a black hole radiation.

[†] The phenomenon called mode conversion has already been known in plasma physics. [6] [7] [8]

[‡]E-mail: saida@phys.h.kyoto-u.ac.jp
[§]E-mail: sakagami@phys.h.kyoto-u.ac.jp

In section II, we explain the MC model [3]. Section III is devoted to the calculation of the spectrum. Lastly, we summarize the results in section IV.

Throughout this paper, we use Planck units, $\hbar = c = G = 1$, and restrict our discussion in two dimension for simplicity.

II. MODE CONVERSION (MC) MODEL

We consider the static spacetime with the metric of the form [3] [4] [5]

$$ds^2 = -dt^2 + (\, dx - v(x)\, dt \,)^2 \,. \tag{1}$$

It is obvious that the timelike Killing vector field of this spacetime is ∂_t. This spacetime has an event horizon at x_h satisfying $v(x_h) = -1$. The surface gravity of this spacetime, κ, is calculated to be $\kappa = (dv/dx)(x_h)$ which is $1/4M$ for the case of the Schwarzschild black hole of mass M. Note that the world line of $dx - v(x)\, dt = 0$ is normal to the surface, $t = \text{const}$, and that it is a geodesic of a free fall observer. The velocity of the free fall observer with respect to the rest observer is $v(x)$, where we require $v(x) < 0$ and monotone increasing $\partial_x v(x) > 0$, that is, $v(x \to \infty) \to v_0$ where $-1 < v_0 \leq 0$ by definition. The proper time of this free fall observer is t of the coordinate (1).

The action of a massless scalar field [3] [4] [5] is

$$S = \frac{1}{2} \int d^2x \sqrt{-\mathrm{g}} \left[(\mathbf{u}(\Phi))^* \, \mathbf{u}(\Phi) - (\mathbf{s}(\Phi))^* \, \mathbf{s}(\Phi) \right] \,, \tag{2}$$

where $\mathrm{g} = \det g_{\mu\nu}$, $\mathbf{u}$ is a unit tangent to the free fall world line and $\mathbf{s}$ is a unit space-like vector which is outward-pointing and orthogonal to $\mathbf{u}$. We choose the derivatives of Φ along $\mathbf{u}$ and $\mathbf{s}$ as $\mathbf{u}(\Phi) = u^\mu \partial_\mu \Phi$ and $\mathbf{s}(\Phi) = \hat{F}(s^\mu \partial_\mu)\Phi$, where $\hat{F}(s^\mu \partial_\mu)$ is a function of the differential operators, ∂_μ, and we require $\hat{F}(-s^\mu \partial_\mu) = -\hat{F}(s^\mu \partial_\mu)$. This modification, $\hat{F}(s^\mu \partial_\mu)$, expresses the MC model's assumption about the effects of unknown Planck scale physics. This leads the EOM of Φ in the coordinate of (1) to be

$$(\partial_t + v(x)\, \partial_x)\, (\partial_t + \partial_x \, v(x))\, \Phi = \hat{F}^2(\partial_x)\Phi \,. \tag{3}$$

Hereafter we specify the form of $\hat{F}(\partial_x)$ as $\hat{F}^2(\partial_x) = \partial_x^2 + \alpha\, \partial_x^4$, where $\alpha = 1/k_c^2$ and k_c is the cut off scale which characterizes the new physics in Planck scale. We set the order of α is of unity, $O(\alpha) \sim 1$.

We require that, at least far from the event horizon, $v(x)$ is nearly constant enough to require the validity of WKB approximation. This means that $\partial_x v(x)/k(x) \ll 1$ and $\partial_x k(x)/k^2(x) \ll 1 \, (k \simeq \text{const.})$ far from the event horizon. With setting $\Phi \simeq \exp(-i\omega t + ikx)$, the EOM (3) gives the dispersion relation

$$\left\{ \begin{array}{lcl} \omega' & = & -vk + \omega \\ \omega' & = & \pm\sqrt{k^2 - \alpha k^4} \equiv F(k) \end{array} \right. , \tag{4}$$

where ω' and ω are the frequencies with respect to the free fall observer and to the rest observer respectively, since $\mathbf{u}(\Phi) = -i(\omega - vk)\Phi = -i\omega'\Phi$ and $\partial_t \Phi = -i\omega\Phi$. The Killing frequency, ω, is conserved during time evolution but the free fall frequency, ω', is not. Equations (4) are the dispersion relation of this MC model, which have four mode solutions for fixed ω. We call the wave numbers of these modes as k_-, k_{-s}, k_{+s} and k_+ in increasing order, as shown in the figure 1. It is not the signature of ω but of ω' according to which we can judge whether the solution is the positive frequency mode or the negative frequency one [3].



FIG. 1. Dispersion relation of our MC model.

The origin of the black hole radiation in this model can be understood by analyzing a wave packet propagation [3] [4] [5]. The group velocity of a wave packet with respect to the free fall observer and the rest observer are expressed as $V_g' = d\omega'(k)/dk$ and $V_g = V_g' + v(x)$ respectively. In the case of $\omega > 0$, we can find easily that $k_{-s,\pm}$ modes are of ingoing, while k_{+s} one is of outgoing. In order to understand the physics of mode conversion, the hydrodynamical analogue of black hole radiation is helpful [2] [3]. In this model, a fluid flows toward a center, on which we consider a wave packet propagation. Here, $v(x)$ in (1) represents the velocity of the infalling fluid flow. As shown in the figure 2, k_{+s} mode travels outward away from the event horizon, and against the infalling flow of back ground fluid. In tracing the mode backward in time, it approaches toward the event horizon. In the figure 3, we show how k_{+s} mode moves in the diagram of the dispersion relation. Near the event horizon, its group velocity, $V_g'(k_{+s})$, eventually coincides with the fluid velocity, $v(x)$. Here the mode conversion takes place. In the view of our tracing backward in time, the infalling wave packet of k_{+s} never reaches the event horizon and is converted to the other wave packet of $k_{-s,\pm}$. We should note that the negative free fall frequency mode of k_- comes to arise in the process of the mode conversion, while the other modes of $k_{-s,+}$ are positive ones.



FIG. 2. Schematical description of the mode conversion on an asymptotically flat spacetime. $\mathrm{I}^{\pm}$ are the future and past null infinity. i_0 is the spatial infinity. H^{+} is the event horizon.



FIG. 3. Graphical description of the mode conversion in the graph of dispersion relation. In view of tracing the wave packet of k_{+s} backward in time, it is converted to the wave packet of $k_{-s,\pm}$.

The situation we consider in the figure 2, where no transmitted wave acrosses the event horizon, corresponds just to a boundary condition for the calculation of the original Hawking radiation. In the asymptotically flat region, $x \to \infty$, we can decompose the wave function as

$$\Phi(x\to\infty) = \sum_{l=\pm s,\pm} c_l \exp(-i\omega + k_l x)\,, \tag{5}$$

where each coefficient, $c_{\pm s,\pm}$, can be uniquely determined by this boundary condition. Then we can obtain the form of the black hole radiation spectrum observed at the asymptotically flat region, $N(\omega)$, by Bogoliubov transformation [3]. It becomes

$$\begin{aligned} N(\omega) &= \langle 0_i|\, a_f^{\dagger} a_f \,|0_i\rangle \\ &= \left| \frac{\omega'(k_-) V_g(k_-) c_-^2(\omega)}{\omega'(k_{+s}) V_g(k_{+s}) c_{+s}^2(\omega)} \right| \,, \end{aligned} \tag{6}$$

where $|0_i\rangle$ is the vacuum state of the free fall observer at the initial time $(t\to-\infty)$, $a_f^{\dagger}$ and a_f are the creation and annihilation operators of free fall observer at the final time $(t \to \infty)$. In deriving (6), we have assumed the Killing frequency spectrum of the wave packet, $\omega(k)$, is sharply peaked around the four wave numbers $k_{\pm s,\pm}$.

III. BH RADIATION SPECRTUM IN MC MODEL

A. Analytical calculation of the spectrum

1. The plan to calculate $N(\omega)$

We consider only the mode function corresponding to the wave number at the peak of $\omega(k)$. We set its form as

$$\Phi(t,x) = \exp(-i\omega t)\,\phi(x)\,, \tag{7}$$

where ω is the peak value of the Killing frequency spectrum of the wave packet. Then EOM (3) becomes

$$\begin{aligned} &\alpha\phi^{(4)} + (1-v^2)\phi'' + \\ &\qquad 2v(i\omega - v')\phi' + (\omega^2 + i\omega v')\phi = 0\,. \end{aligned} \tag{8}$$

As mentioned at the end of previous section, we adopt the boundary condition of total reflection, which is expressed as

B.C.: *The solution of* (8) *dumps with decreasing x inside the event horizon.*

We take the origin of spatial coordinate x at the location of the event horizon, that is, $v(x_h = 0) = -1$.

To calculate $N(\omega)$, we follow the same plan as in [4]. This plan consists of three steps: First step is to solve (8) around the event horizon. We use the Laplace transformation in this step, where the contour of the Laplace integral should be chosen as it satisfies the B.C. Secondly, we look for the solution of (8) far from the event horizon by means of the WKB approximation. We match the WKB solutions with those of the first step. The existence of the overlap region (matching zone) should be checked, where the solutions in first and second steps are valid. Third step is to calculate $N(\omega)$ at the spatial infinity with using (6).

2. Formulas for each step and small parameters

We expand $v(x)$ around the event horizon as

$$\begin{aligned} v(x) &\simeq -1 + \kappa x\,, \\ 1 - v(x)^2 &\simeq 2\kappa x\,, \end{aligned} \tag{9}$$

where we require $|\kappa x| \ll 1$ and κ is the surface gravity which is assumed to be small itself. We use Laplace transformation of $\phi(x)$,

$$\phi(x) = \int_C ds\, e^{sx}\, \tilde{\phi}(s)\,. \tag{10}$$

With approximating every coefficient in EOM (8) up to $O(\kappa)$, we obtain

$$2\kappa f(s)\,\tilde{\phi}'(s) - B(s)\,\tilde{\phi}(s) = 0\,, \tag{11}$$

where $f(s) = s^2 + i\omega s$, $B(s) = \alpha s^4 - \kappa\beta f'(s)$ and $\beta = 1 + i\omega/\kappa$. This gives us

$$\begin{aligned}
&\phi(x) \\
&= \int_C ds\ (s+i\omega)^{i\alpha\omega^3/2\kappa}\left(s^2+i\omega s\right)^{-\beta/2} \times \\
&\qquad \exp\left[sx + \frac{\alpha}{2\kappa}\left(\frac{1}{3}s^3 - i\frac{\omega}{2}s^2 - \omega^2 s - i\frac{11}{6}\omega^3\right)\right] \\
&\equiv \int_C ds\ \exp\left(h(s)\right)\,.
\end{aligned} \tag{12}$$

We will find later that the factor $(s+i\omega)^{i\alpha\omega^3/2\kappa}$, which is ignored in [4], will make important contributions to the deviation of the MC spectrum from the thermal one.

For WKB approximation, we set

$$\phi(x) = \exp\left[i\int_{x_0}^{x} k(x')dx'\right]\,, \tag{13}$$

where x_0 is a constant. We expand $k(x) = k_0(x) + (1/\lambda)k_1(x) + \cdots$ in introducing the scaling $x \to \lambda x$ $(\partial_x \to (1/\lambda)\partial_x)$ [4], then we obtain from (8) that

$$O(1) : \alpha k_0^4 - (1-v^2)k_0^2 - 2\omega v k_0 + \omega^2 = 0 \tag{14}$$

$$\begin{aligned}
O(\lambda^{-1}) : k_1 = \frac{i}{2}\frac{d}{dx}\ln\big[\sqrt{\alpha}\,\big(2\alpha k_0^3 \\
-(1-v^2)k_0 - \omega v\big)\big]\,,
\end{aligned} \tag{15}$$

where four solutions of k correspond to $k_{\pm,\pm s}$.

To calculate $N(\omega)$, we should evaluate $\omega' V_g$. This is given by (4) to be

$$\omega' V_g = -2\alpha k_0^3 + (1-v_0^2)k_0 + \omega v_0\,. \tag{16}$$

As stated in section I, $N(\omega)$ deviates from the thermal spectrum in the high frequency region $\kappa < \omega$ [3]. So we may not be able to derive the deviation with expanding k_0 around $\omega = 0$ as done in [4]. Thus we focus our attention on the region, $\kappa < \omega$, in order to derive the deviation. It is expected that the deviation becomes visible at the value of ω which does not exceed κ so much. Thus, when we analyze how the MC spectrum deviates from the thermal one, the fourth term in the left-hand-side of (14) can be neglected in obtaining WKB solutions. This means to require

$$\alpha k_0^4 \gg \omega^2\,,\ (1-v^2)k_0^2 \gg \omega^2\,,\ \omega v k_0 \gg \omega^2\,. \tag{17}$$

The other restrictions on the region of κ and ω come from the validity of the solution of WKB and that of the Laplace transformation. These restrictions are expressed by $|k_1/k_0| \ll 1$ from WKB and by $\kappa x \ll 1$ and $|s_0 x| \gg 1$ from Laplace transformation, where s_0 is a saddle point of integrand in (12) in using the asymptotic expansion of x by the method of steepest descent contour (SDC). We can obtain the matching zone from these inequalities. It becomes

$$\left(\frac{\alpha}{\kappa}\right)^{1/3} \ll x \ll \frac{1}{\kappa}\,, \tag{18}$$

where we use the forms of s_0 and $k_{0,1}$ which are calculated later in appendix A and at (23). Furthermore, we require $\omega x \ll 1$ for later convinience.

With introducing λ and μ by $\kappa x = (1/\kappa\sqrt{\alpha})^\lambda$ and $\omega = \kappa(1/\kappa\sqrt{\alpha})^\mu$ where $\kappa\sqrt{\alpha} \ll 1$, the parameter region we discussed above can be expressed as

$$3\lambda > 2(\mu - 1)\,,\ -\frac{2}{3} < \lambda < 0\,,\ 0 < \mu\,,\ \lambda + \mu < 0\,, \tag{19}$$

where first and second inequalities obtained by (17) with (9) and (18) respectively.



FIG. 4. Parameter region we consider.

3. *Calculation of* $N(\omega)$

We can proceed to calculate $N(\omega)$ with above preparations. As we carry our the Laplace integral of (12) in the appendix A, we obtain the solution of (8) in the matching zone as

$$\phi(x) = \phi_1(x) + \phi_2(x) + \phi_3(x)\,, \tag{20}$$

where, as demonstrated below, $\phi_{1,2,3}$ should be matched with WKB solutions, $\phi_{\pm,+s}$, respectively. These are expressed as

$$\begin{aligned}
&\phi_{1,2}(x) \\
&= \sqrt{\pi}\,e^{i\pi(n+1/2-\varepsilon 3/4)-i11\alpha\omega^3/12\kappa} \times \\
&\qquad e^{\varepsilon\pi\omega/2\kappa(1-\alpha\omega^2/2)}\left(\frac{2\kappa}{\alpha}\right)^{-1/4-i\omega/2\kappa+i\alpha\omega^3/4\kappa} \times \\
&\qquad x^{-3/4-i\omega/2\kappa}\ \exp\left[\varepsilon i\frac{2}{3}\sqrt{\frac{2\kappa}{\alpha}}x^{3/2} + i\frac{\omega}{2}x\right]\,,
\end{aligned} \tag{21}$$

$$\begin{aligned}\phi_3(x)& \\ =\ & e^{-5\xi/3}\,2\,\sinh\left[\frac{\pi\omega}{\kappa}\left(1-\frac{\alpha\omega^2}{2}\right)\right]\times \\ & \Gamma\left[-i\frac{\omega}{\kappa}\left(1-\frac{\alpha\omega^2}{2}\right)\right]x^{i\omega/\kappa}\,,\end{aligned} \qquad (22)$$

where $(\varepsilon, n) = (-1, 0)$, $(1, 1)$ for ϕ_1 and ϕ_2 respectively and $\xi = i\alpha\omega^3/4\kappa$. The factor $1-\alpha\omega^2/2$ shows the difference form the analysis in [4], which arises from the factor $(s+i\omega)^{i\alpha\omega^3/2\kappa}$ in the integrand of (12).

Next we shift our step to obtaining WKB solutions. We can obtain k_0 from (14) with the conditions of (17) only. Then k_0 becmoes

$$k_0 \simeq \begin{cases}\omega/(1+v) & \text{for } k_{+s} \\ \pm\sqrt{(1-v^2)/\alpha}+v\omega/(1-v^2) & \text{for } k_\pm\end{cases}\,, \qquad (23)$$

where these forms of k_0 are valid in the region $3\lambda > 2(\mu-1)$ which is given by (17). Thus, for the purpose of matching with (20), we obtain WKB solutions in the parameter region of (19) as

$$\begin{aligned}\phi_\pm(x) &\simeq A_\pm\,(\pm 2\kappa)^{-3/4}\,x^{-3/4-i\omega/2\kappa}\times \\ & \exp\left[\pm\frac{2}{3}\sqrt{\frac{2\kappa}{\alpha}}x^{3/2}+i\frac{\omega}{2}x\right]\,, \qquad (24)\\ \phi_{+s}(x) &\simeq A_{+s}x^{i\omega/\kappa}\,, \qquad (25)\end{aligned}$$

where $A_{\pm,+s}$ are the coefficients obtained by x_0 in (13).

By matching WKB solutions, (24) and (25), with that of Laplace transformation, (20), we obtain the wave function as

$$\begin{aligned}\phi(x)& \\ =\ & \sqrt{\pi}\,e^{i9\pi/4-11\xi/3}\times \\ & e^{-\pi\omega/2\kappa(1-\alpha\omega^2/2)}\left(\frac{2\kappa}{\alpha}\right)^{-1/4-i\omega/2\kappa+\xi}\times \\ & \left[\frac{(-2\kappa)^{3/4}}{A_-}\,\phi_-(x)\right. \\ & \left.\quad+e^{-i5\pi/2+\pi\omega/\kappa(1-\alpha\omega^2/2)}\,\frac{(2\kappa)^{3/4}}{A_+}\,\phi_+(x)\right] \\ +\ & e^{-5\xi/3}\,2\,\sinh\left[\frac{\pi\omega}{\kappa}\left(1-\frac{\alpha\omega^2}{2}\right)\right]\times \\ & \Gamma\left[-i\frac{\omega}{\kappa}\left(1-\frac{\alpha\omega^2}{2}\right)\right]\frac{1}{A_{+s}}\,\phi_{+s}(x)\,.\end{aligned} \qquad (26)$$

Note that we can use this result beyond the matching zone, i.e. $(\alpha/\kappa)^{1/3} \ll x$ which is given by the validity of WKB approximaton, $|k_1/k_0| \ll 1$.

To obtain $N(\omega)$, we need to evaluate $\omega' V_g$ and $\phi(x)$ at the asymptotically flat region. We obtain with taking the limit $v(x\to\infty) \to v_0$ that

$$\begin{aligned}\phi_\pm(x\to\infty) &\simeq A_\pm\,(\pm 1)^{-1/2}(1-v_0^2)^{-3/4}\times \\ & \times\exp\left[\pm i\sqrt{\frac{1-v_0^2}{\alpha}}x+i\frac{v_0\omega}{1-v_0^2}x\right]\,, \qquad (27) \\ \phi_{+s}(x\to\infty) &\simeq A_{+s}\exp\left[-i\frac{2v_0\omega}{1-v_0^2}x\right]\,, \qquad (28)\end{aligned}$$

and

$$\begin{aligned}\omega'(k_\pm)V_g(k_\pm) &\simeq \mp\frac{(1-v_0^2)^{3/2}}{\sqrt{\alpha}}\,, \qquad (29) \\ \omega'(k_{+s})V_g(k_{+s}) &\simeq -\omega\,. \qquad (30)\end{aligned}$$

Then we get c_- and c_{+s} by (26), (27) and (28), and finally obtain $N(\omega)$ by (6) to be

$$N(\omega) = \frac{1-\alpha\omega^2/2}{\exp[(2\pi\omega/\kappa)\,(1-\alpha\omega^2/2)]-1}\,. \qquad (31)$$

This result shows that MC spectrum is enhanced in comparison with Hawking spectrum.

B. Comparison with numerical results

We compare the analytically derived form of the spectrum, (31), with numerical calculations. The numerical calculations are carried out with two types of $v(x)$, which are shown successively in the following two subsections.

We have used *Mathematica* to solve (8) and fit its solution with the form, $\phi = \sum_{l=\pm,\pm S} c_l \exp(ik_l)$, in order to get c_- and c_{+s}.

Note that we have set $\kappa/2\pi = 0.0008$ and $\alpha = 1$ througout these numerical computations.

1. First type of $v(x)$ – CJ type

First example of $v(x)$, which we call Corley-Jacobson (CJ) type [3], takes the form

$$v(x) = \mathrm{sgn}(x)\,\frac{1}{2}\sqrt{\tanh\left[(2\kappa x)^2\right]}-1\,. \qquad (32)$$

To let k_- mode be a negative free fall frequency one at the asymptotically flat region, the value of ω is restricted by upper value $\omega_{max} \sim 0.16$. Further, in our parameter region (19), the range of ω is bounded as $0.005 < \omega < 0.04$ $(0 < \mu < 2/5)$.

We have calculated the relative deviation,

$$R(\omega) = \frac{N(\omega)}{N_H(\omega)}-1\,, \qquad (33)$$

where N_H is the Hawking spectrum. Figure 5 is the plot of this relative deviation. $R(\omega)$ grows qualitatively more and more with increasing ω.



FIG. 5. Graph of relative deviation, $R(\omega)$, with CJ type of $v(x)$. $R(\omega)$ grows qualitatively more and more with increasing ω.

This shows that the MC spectrum is enhanced in comparison with the Hawking spectrum, as is indicated by the analytical result (31). However figure 8, which is the plots of the logarithm of relative deviations, exhibit that the order of $R(\omega)$ in CJ type differs largely from our result (31). We expect that this difference can be reduced by changing the form of $v(x)$. The clue of this change is in the EOM (8). The most effective term for determining the analyticity of the solution of (8) is the second term, $(1-v^2)\phi''$. [**] Thus, in order to compare (31) with a numerical calculation, the expansion $1-v^2 \simeq 2\kappa x$ near horizon should be better than the expansion $v \simeq -1+\kappa x$. By the way, $v(x)$ of CJ type can be expanded near horizon as:

$$\begin{aligned} v(x) &\simeq -1+\kappa x + O\left((\kappa x)^4\right) , \\ 1-v(x)^2 &\simeq 2\kappa x + O\left((\kappa x)^2\right) . \end{aligned} \tag{34}$$

While CJ type is good form for the expansion $v \simeq -1+\kappa x$ near horizon, however it is not good one for the expansion $1-v^2 \simeq 2\kappa x$. We should modify $v(x)$ to let the expansion $1-v^2 \simeq 2\kappa x$ near horizon be better than CJ type.

2. Second type of $v(x)$ – DS type

The modified type of $v(x)$, which is called as Double-Squareroot-Tanh (DS) type hereafter, is

$$v(x) = -\sqrt{1-\mathrm{sgn}(x)\frac{3}{4}\sqrt{\tanh\left[\left(\frac{8}{3}\kappa x\right)^2\right]}} . \tag{35}$$

The range of ω is $0.005 < \omega < 0.04$ $(0 < \mu < 2/5)$, because of the same reason as that of CJ type. Figure 6 shows the shapes of $v(x)$ of CJ and DS types. They vary from -1 at $x=0$ to $x=-1/2$ at $x \to \infty$.



FIG. 6. Shapes of $v(x)$ in the form of CJ and DS types. They vary from -1 at $x=0$ to $x=-1/2$ at $x \to \infty$.

The $v(x)$ of DS type takes the form near horizon:

$$\begin{aligned} v(x) &\simeq -1+\kappa x + O\left((\kappa x)^2\right) , \\ 1-v(x)^2 &\simeq 2\kappa x + O\left((\kappa x)^5\right) . \end{aligned} \tag{36}$$

The expansion $v \simeq -1+\kappa x$ near horizon for CJ type is better than that for DS type, however the expansion $1-v^2 \simeq 2\kappa x$ for DS type is better than that for CJ type.

Figure 7 is the plot of the relative deviation for DS type. $R(\omega)$ grows qualitatively more and more with increasing ω. This figure and the figure 5 implies that the details of the MC spectrum depend strongly on the form of $v(x)$. The total behavior of the figure 7, however, shows that, althogh some data take negative value, the qualitative nature of the MC spectrum is enhancement in comparison with the Hawking spectrum, which is indecated by the analytical result (31). Then the figure 8 involves the plot of the absolute value of $R(\omega)$ in DS type. This shows that the order of $R(\omega)$ can be explained by our analytical result (31) fairly good, since the higher order effects by $1-v^2$ in DS type are weakened to agree with the assumption made in our analytical treatment of the MC spectrum.

From above it is valid to mention that our resultant spectrum (31) can explain the qualitative nature of the MC spectrum in the range, $\kappa < \omega < \kappa(\kappa\sqrt{\alpha})^{-2/5}$.

[**] This is also implied in section IV.



FIG. 7. Graph of relative deviation, $R(\omega)$, with DS type of $v(x)$. $R(\omega)$ grows qualitatively more and more with increasing ω.



FIG. 8. Graphs of logarithm of absolute value of the relative deviations. The solid line is for our analytical result. These show that our resultant spectrum explains the qualitative nature of the MC spectrum.

IV. SUMMARY AND DISCUSSION

We have derived the correction to the black hole radiation spectrum in the MC model in the frequecy range, $0 < \mu < 2/5$, $\omega = \kappa(1/\kappa\sqrt{\alpha})^{\mu}$. The MC model cut the high frequency mode off, so one may guess naively that the spectrum would be suppressed. Our result, however, shows that the MC spectrum is enhanced in the frequency range, $0 < \mu < 2/5$. This is consistent with the numerical calculations as seen above.

The deviation factor $1-\alpha\omega^2/2$ in (31) is understood as a correction of red shift which outgoing modes undergo during propagating from the event horizon to the asymptotically flat region. This is reflected on the form of k_{+s} mode, $x^{i(\omega/\kappa)(1-\alpha\omega^2/2)}$ in (A6). This form can be easily explained. Neglecting the fourth term in (8) which is of second order of small quantities κ and ω, we obtain

$$\alpha\varphi''' + 2\kappa x\varphi' - 2(1-\kappa x)(i\omega - \kappa)\varphi = 0\,, \tag{37}$$

where we set $\phi'(x) = \varphi(x)$ and $v(x) \simeq -1 + \kappa x$. Under the assumption $\alpha < 1$, we expand $\varphi = \varphi_0 + \alpha\varphi_1 + \cdots$ which leads

$$2\kappa x\varphi_0' - 2(1-\kappa x)(i\omega - \kappa)\varphi_0 = 0 \tag{38}$$
$$2\kappa x\varphi_1' - 2(1-\kappa x)(i\omega - \kappa)\varphi_1 = -\varphi_0'''\,. \tag{39}$$

The lowest order solution is given by

$$\varphi_0(x) = x^{i\omega/\kappa - 1} e^{(\kappa - i\omega)x}\,. \tag{40}$$

We are interested in the analyticity of the following form:

$$\varphi_0 \sim x^{i\omega/\kappa - 1} \quad \Rightarrow \quad \phi_0 \sim x^{i\omega/\kappa}\,, \tag{41}$$

since the temperature of the original Hawking radiation is determined by the form of an outgoing mode which takes the same form as above. Then we consider the correction to this analyticity. With setting $B = i\omega/\kappa - 1$, we obtain φ_1 as

$$\begin{aligned} &\varphi_1(x) \\ &= -\frac{\alpha}{2}\left[-\frac{B^3 - 3B^2 + 2B}{3\kappa x^3} + \frac{3(B^3 - B^2)}{2x^2}\right. \\ &\qquad \left. -\frac{3\kappa B^3}{x} - \kappa^2 B^3 \ln x\right]\varphi_0(x)\,. \end{aligned} \tag{42}$$

We can recognize the fourth term in the square brackets as the secular one. Although this term is not dominant, it can change the analyticity which determines the temperature of the radiation. We use the renormalization group method in order to take all of the contributions of the secular term of φ_1 into account [9] [10]. In the parameter region $\omega/\kappa \geq 1$, we obtain

$$\begin{aligned} &\varphi \sim x^{\alpha(i\omega-\kappa)^3/2\kappa + i\omega/\kappa - 1} \sim x^{\alpha(i\omega)^3/2\kappa + i\omega/\kappa - 1} \\ &\Rightarrow \phi \sim x^{i(\omega/\kappa)(1-\alpha\omega^2/2)}\,, \end{aligned} \tag{43}$$

which shows the same deviation as in (31).

Further, our result agrees with that of [11]. In the paper [11], the order of the MC spectrum's deviation has been estimated, which is consistent with our spectrum, $N(\omega)$.

ACKNOWLEDGEMENTS

We would like to Thank J.Soda for his useful discussions. This work was supported in part by Monbusho Grant-in-Aid for Scientific Reserch No. 10640258. One of us (H.S.) also thanks S.Kawai for his helpful comments.

APPENDIX A: CALCULATION OF LAPLACE INTEGRAL

We approximate (12) in this appendix by an asymptotic expansion of x with the method of steepest descent contour (SDC). The saddle points, s_0, are the solutions of $dh/ds = 0$:

$$\alpha s_0^4 + 2\kappa x s_0^2 + 2\left(i\omega\kappa x - \kappa\beta\right) s_0 - i\omega\kappa\beta = 0\,. \tag{A1}$$

The $\phi(x)$ approximated by SDC is expressed as

$$\phi(x) \simeq \sqrt{2\pi}\,\left(|x| h''(s_0)\right)^{-1/2} \times \exp\left[x h(s_0) + i\pi\left(\frac{1}{2}\gamma + n\right)\right]\,, \tag{A2}$$

where n is the integer determining the direction of the SDC and $\gamma = 0,\, 1$ for $x < 0$ and $x > 0$ respectively.

With approximating (A1) and (A2) in the paramater region (19), we obtain the saddle point satisfying B.C. in $x < 0$ as

$$s_0 \simeq \sqrt{2\kappa|x|/\alpha} - i\beta/2|x|\,, \tag{A3}$$

and for $x > 0$, the saddle points are

$$s_0 \simeq \begin{cases} \beta/x \\ \pm i\sqrt{2\kappa x/\alpha} - \beta/2x \end{cases}. \tag{A4}$$

The integral contour is determined as shown in the figure 9, where the contour should asymtpote to one of the three regions, $\pi/6 < \arg(s) < \pi/2$, $5\pi/6 < \arg(s) < 7\pi/6$ and $3\pi/2 < \arg(s) < 11\pi/6$ since, in the limit $|s| \to \infty$, the convergence of (12) requires $Re(s^3) < 0$. The waving curves are the branch cuts of the integrand in (12).



FIG. 9. Contour of Laplace integral satisfying B.C. Left figure is for $x < 0$. Right figure is for $x > 0$.

The SDC of saddle point β/x approximates the contribution from the contour C_3. We can find, however, that this contribuiton is obtained by integral representations of some special functions more correctly than by the SDC approxiamtion. So we use SDC approximation only for the contributions from the contours C_1 and C_2. The dominant dependence on x of these contributions are obtained by (A2) to give us (21).

To carry out the contour integral of C_3, we transform s to t by $sx = e^{-i\pi}t$. While we choose the argument of s as $-\pi < \arg(s) < \pi$, the argument of t is to be $0 < \arg(t) < 2\pi$. Then (12) becomes

$$\begin{aligned} &\phi_3(x) \\ &= e^{-i\pi - i11\alpha\omega^3/12\kappa}\,(i\omega)^{-\beta/2 + i\alpha\omega^3/2\kappa}\,x^{\beta/2-1} \times \\ &\quad \int_{C_t} dt\,(-t)^{-\beta/2}\left(1 + \frac{t}{-i\omega x}\right)^{-\beta/2 + i\alpha\omega^3/2\kappa} \times \\ &\quad \exp\left[-t + \frac{\alpha}{2\kappa}\left(-\frac{t^3}{3x^3} - i\frac{\omega t^2}{2x^2} + \frac{\omega^2 t}{x}\right)\right]\,, \end{aligned} \tag{A5}$$

where C_t is the contour on t-plane corresponding to C_3, and $i\omega = e^{i\pi/2}\omega$ and $-i\omega x = e^{i3\pi/2}\omega x$. We can simplify the exponent of (A5) to be $\exp(-t)$ in our parameter region (19) by the same argument in [4]: by $1/|\kappa x^3| \ll 1$ and $\omega x \ll 1$, the correction by $\exp[(\alpha/2\kappa)(\cdots)] \simeq 1 + (\alpha/2\kappa)(\cdots)$ is negligible. With dividing the contour C_t into C_A and C_B as shown in the figure 10 and using the integral representation of Whittaker function, we can express (A5) as the sum of two Whittaker functions.



FIG. 10. Contour of Laplace integral for $\phi_3(x)$.

Further we can simplify it by one confluent hypergeometric function to be

$$\begin{aligned} &\phi_3(x) \\ &= e^{-11\xi/3}\,2\,\sinh\left[\frac{\pi\omega}{\kappa}\left(1 - \frac{\alpha\omega^2}{2}\right)\right] \times \\ &\quad \Gamma\left[-i\frac{\omega}{\kappa}\left(1 - \frac{\alpha\omega^2}{2}\right)\right] \times \\ &\quad e^{-i\omega x}\,x^{i\omega/\kappa(1-\alpha\omega^2/2)} \times \\ &\quad F\left[\frac{\beta}{2}, 1 + i\frac{\omega}{\kappa}\left(1 - \frac{\alpha\omega^2}{2}\right); i\omega x\right]\,, \end{aligned} \tag{A6}$$

where $\xi = i\alpha\omega^3/4\kappa$, $\beta = 1 + i\omega/\kappa$, Γ is the gamma function and F is the confluent hypergeometric function. Then we obtain the dominant dependence on x of ϕ_3, which bocomes (22).

---

[1] S.W.Hawking, Commun.math.phys.**43** 199(1975)

[2] W.G.Unruh, Phys.Rev.D**51** 2827(1995)

[3] S.Corley and T.Jacobson, Phys.Rev.D**54** 1568(1996), hep-th/9601073

[4] S.Corley, Phys.Rev.D**57** 6280(1998), hep-th/9710075

[5] S.Corley and T.Jacobson, Phys.Rev.D**57** 6269(1998), hep-th/9709166

[6] D.G.Swanson, *Plasma Waves* (Academic Press, 1989)

[7] T.H.Stix, *Waves in Plasmas* (American Institute of Physics, 1992)

[8] D.G.Swanson, *Theory of Mode Conversion and Tunneling in Inhomobeneous Plasmas*, (John Wiley & Sons, New York, 1998)

[9] L.Y.Chen, N.Goldenfeld and Y.Oono, Phys.Rev.Lett.**73** 1311(1994)

[10] L.Y.Chen, N.Goldenfeld and Y.Oono, Phys.Rev.E**54** 376(1996), hep-th/9506161

[11] Y.Himemoto and T.Tanaka, gr-qc/9904076